



Important Safety Information

Do not use LATISSE™ solution if you are allergic to one of its ingredients.

LATISSE™ solution is intended for **use on the skin of the upper eyelid margins at the base of the eyelashes. DO NOT APPLY** to the lower eyelid. If you have a history of abnormal IOP, you should only use LATISSE™ under the close supervision of your physician.

LATISSE™ use may cause darkening of the eyelid skin which may be reversible.

You should tell your physician you are using LATISSE™ especially if you have a history of eye pressure problems. You should also tell anyone conducting an eye pressure screening that you are using LATISSE™.

The most common side effects after using LATISSE™ solution are an itching sensation in the eyes and/or eye redness. This was reported in approximately 4% of patients. LATISSE™ solution may cause other less common side effects which typically occur on the skin close to where LATISSE™ is applied, or in the eyes. These include skin darkening, eye irritation, dryness of the eyes, and redness of the eyelids.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Be advised that as an alternative to purchasing Latisse at Commonwealth Facial Plastic surgery, you can be provided a prescription that can be filled at a pharmacy.

I understand that this product is non-returnable. ALL SALES ARE FINAL

Patient _____

Date _____

Witness _____

Date _____